

An Extension of the Fluctuation Theorem

R. van Zon and E.G.D. Cohen

The Rockefeller University
1230 York Avenue, New York, NY 10021

May 7, 2003

Abstract

Heat fluctuations are studied in a dissipative system with both mechanical *and* stochastic components for a simple model: a Brownian particle dragged through water by a moving potential. An extended stationary state fluctuation theorem is derived. For infinite time, this reduces to the conventional fluctuation theorem only for small fluctuations; for large fluctuations, it gives a much larger ratio of the probabilities of the particle to absorb rather than supply heat. This persists for finite times and should be observable in experiments similar to a recent one of Wang *et al.*

PACS: 05.40.-a, 05.70.-a, 44.05.+e, 02.50.-r

1 Introduction

There is a lack of unifying principles in nonequilibrium statistical mechanics, compared to the equilibrium case. So it is not surprising that the *fluctuation theorem* has received a lot of attention, as it gives a property of fluctuations of entropy production for a large class of systems, possibly arbitrarily far from equilibrium[1, 2, 3, 4, 5, 6, 7, 8, 9, 10].

In terms of heat rather than entropy production, the conventional stationary state fluctuation theorem (SSFT)¹ states that the probability $P_\tau(Q_\tau)$

¹We call equation (1) the “conventional SSFT” to distinguish it from the extended SSFT presented in this paper.

to find a value of Q_τ for the amount of heat dissipated in a time interval τ , satisfies, in a nonequilibrium stationary state[1, 3],

$$\frac{P_\tau(Q_\tau)}{P_\tau(-Q_\tau)} \sim e^{\beta Q_\tau}, \quad (1)$$

where $\sim$ indicates the behavior for large τ . Here $\beta = (k_B T)^{-1}$, with k_B Boltzmann's constant and T the (effective) temperature of the system. In contrast, the transient fluctuation theorem (TFT) considers fluctuations Q_τ in time, when the system is initially in equilibrium[2]. These theorems were first demonstrated in deterministic many particle systems in an external field [1, 2, 3], but later in stochastic systems as well[4, 5].

In this context, it is customary to identify Q_τ/T as a (generalized) entropy production. Then, equation (1) is interpreted as a theorem for entropy production arbitrarily far from equilibrium. Furthermore, the fluctuation theorem holds for arbitrary values of Q_τ , i.e., also far from its average. Hence, it is often referred to as a large deviation theorem.

Recently, a laboratory experiment was carried out by Wang *et al.*[8]. They measured fluctuations in the work done on a system in a transient state of a Brownian particle in water, subject to a moving, confining potential. The TFT was confirmed.

In the deterministic models, dissipation is often modeled by including a damping term in the equations of motion, chosen such as to keep, e.g., the total energy of the system fixed. This damping term is a mechanical expression for what in reality is an external thermostat. Any work done by the external field on the system is absorbed by this “internal” thermostat. So the external work done on the system equals the heat dissipated in the system.

While Wang *et al.* intended to study the entropy production (or heat) fluctuations, in fact, the work fluctuations were studied[10]. In contrast to the above sketched purely deterministic models, work fluctuations differ from heat fluctuations in their system due to the presence of a confining potential. Thus, some of the external work done is converted into potential energy and only the rest is converted into heat. In fact, in this system the work fluctuations in the stationary state satisfy the conventional SSFT[10, 11], but the heat fluctuations do not, as we shall show. As it turns out, having a *deterministic* together with a *stochastic* motion results in a behavior of the heat fluctuations that coincides with the conventional SSFT only for a

restricted set of small fluctuations and in a very different behavior for larger ones.

2 Work fluctuations – conventional SSFT

We first discuss the work-related fluctuation theorem in the experiment of Wang *et al.*, which was treated theoretically in Refs. [10, 11, 12]. The Brownian particle in a fluid, subject to a harmonic potential moving with constant velocity $\mathbf{v}^*$, was described by an overdamped Langevin equation:

$$\frac{d\mathbf{x}_t}{dt} = -(\mathbf{x}_t - \mathbf{x}_t^*) + \zeta_t. \quad (2)$$

Here, $\mathbf{x}_t$ is the position of the particle at time t , $\mathbf{x}_t^* = \mathbf{v}^*t$ is the position of the minimum of the harmonic potential at time t [cf. equation (7')], and ζ_t is a fluctuating force with zero mean and a delta function correlation in time. We remark that the relaxation time of the position of the particle has been set equal to one. Also, we set $k_B T = 1$, so $\langle \zeta_t \zeta_s \rangle = 2\delta(t - s)$.² In reference [10], it was shown that equation (2) is solvable in a co-moving frame, in which it reduces to a standard Ornstein-Uhlenbeck process. Thus, the stationary probability distribution and Green's function are known, and are both Gaussian in $\mathbf{x}_t$. The work is the total amount of energy put into the system in a time τ . This is a fluctuating quantity, given by³

$$W_\tau \equiv \mathbf{v}^* \cdot \int_0^\tau [-(\mathbf{x}_t - \mathbf{x}_t^*)] dt. \quad (3)$$

Here, the time $t = 0$ denotes the initial time of an interval of length τ in the stationary state. W_τ is a linear function of the positions $\mathbf{x}_t$, and since those have a Gaussian probability distribution function, so does W_τ . When the mean and variance of the probability distribution function P_τ^W are computed [using the stationary solution and Green's function of equation (2)], one finds[10]

$$\lim_{\tau \rightarrow \infty} \frac{1}{w\tau} \ln \left[\frac{P_\tau^W(pw\tau)}{P_\tau^W(-pw\tau)} \right] = p. \quad (4)$$

²Compared to Ref. [10], $k_B T = 1$, the harmonic force constant $k = 1$ and the friction coefficient $\alpha = 1$. So the energy unit is $k_B T$, the time unit is the relaxation time α/k , and the length unit is the thermal width $\sqrt{k_B T/k}$.

³A force on the particle implies an opposite force on U , which must be overcome by an external force to keep U moving. The work done by this force is W_τ in equation (3)[10].

Here, p is a scaled value of W_τ , defined as $p = W_\tau / \langle W_\tau \rangle$, such that $\langle p \rangle = 1$. We also wrote

$$\langle W_\tau \rangle = w\tau, \quad (5)$$

with w the average work rate, which is independent of τ in the stationary state. In the current units, $w = |\mathbf{v}^*|^2$. Equation (4) is, for the work fluctuations, a more careful formulation of the SSFT in equation (1). A work related TFT also holds[10, 11, 12].

3 Heat fluctuations

3.1 Fourier transform of the distribution

We now turn to the heat SSFT. The heat Q_τ is that part of the work W_τ that goes into the fluid. Some work is also stored in the potential, so

$$Q_\tau \equiv W_\tau - \Delta U_\tau, \quad (6)$$

where ΔU_τ is the change in potential energy of the particle in a time τ ,

$$\Delta U_\tau \equiv U_\tau - U_0, \quad (7)$$

with

$$U_t \equiv \frac{1}{2} |\mathbf{x}_t - \mathbf{x}_t^*|^2. \quad (7')$$

This form of U_t makes Q_τ nonlinear in $\mathbf{x}_t$. As a result, the probability distribution function $P_\tau(Q_\tau)$ of Q_τ need not be Gaussian. Nonetheless, it is possible to compute its Fourier transform.

The Fourier transform of $P_\tau(Q_\tau)$, defined as

$$\hat{P}_\tau(q) \equiv \int_{-\infty}^{\infty} dQ_\tau e^{iqQ_\tau} P_\tau(Q_\tau), \quad (8)$$

is computed by writing P_τ as [using equations (6) and (7)]

$$P_\tau(Q_\tau) = \iint d\mathbf{x}_0 d\mathbf{x}_\tau P_\tau^{W_\tau, \mathbf{x}_0, \mathbf{x}_\tau}(Q_\tau + \Delta U_\tau, \mathbf{x}_0, \mathbf{x}_\tau), \quad (9)$$

where $P_\tau^{W_\tau, \mathbf{x}_0, \mathbf{x}_\tau}$ is the joint distribution of the work W_τ , the positions $\mathbf{x}_0$ and $\mathbf{x}_\tau$ at the beginning and at the end of the time interval τ , respectively.

This distribution is Gaussian because W_τ , $\mathbf{x}_0$, and $\mathbf{x}_\tau$ are all linear in $\mathbf{x}_t$. When equation (9) is inserted into equation (8), a seven dimensional Gaussian integral is left, which after some algebra yields

$$\hat{P}_\tau(q) = \frac{\exp \left\{ wq(i-q) \left[\tau - \frac{2q^2(1-e^{-\tau})^2}{1+(1-e^{-2\tau})q^2} \right] \right\}}{[1 + (1 - e^{-2\tau})q^2]^{3/2}}. \quad (10)$$

Once $\hat{P}_\tau(q)$ has been transformed back, one considers

$$f_\tau(p) \equiv \frac{1}{w\tau} \ln \left[\frac{P_\tau(pw\tau)}{P_\tau(-pw\tau)} \right]. \quad (11)$$

Here, p is a scaled value of Q_τ , defined as

$$p = Q_\tau / \langle Q_\tau \rangle,$$

i.e., $\langle p \rangle = 1$. We also used that

$$\langle Q_\tau \rangle = \langle W_\tau \rangle - \langle \Delta U_\tau \rangle = w\tau$$

by equation (5), since $\langle \Delta U_\tau \rangle = 0$ in the stationary state. The scaled logarithmic ratio $f_\tau(p)$ should be equal to p for $\tau \rightarrow \infty$ when the conventional SSFT holds.

As far as we know, there is no exact result for the inverse Fourier transform of $\hat{P}_\tau(q)$ in equation (10) in terms of known functions. Therefore, a completely analytic treatment did not seem feasible. Instead, we used first a numerical method, the fast Fourier transform algorithm[13], to invert equation (10). The resulting probability distribution function P_τ as well as the corresponding f_τ have been plotted in figure 1. These results do not agree very well with the straight line with slope one, which should be approached for large τ if the conventional SSFT were to hold. One might think that this is due to τ not being large enough. However, we found that deviations of $f_\tau(p)$ from p for large p are generic, while the straight line *is* approached only for smaller p . Nonetheless, we cannot say anything conclusive about the large τ , large p behavior because the distribution gets very peaked and hence becomes smaller for large deviations, which makes the numerical method unreliable.




Figure 1: Numerically obtained $f_\tau(p)$ (bold dashed) for $v^* = 1.5$ and $\tau = 1.3$, versus the (v^* independent) extension of the SSFT for $\tau \rightarrow \infty$ (bold solid). Also plotted are the conventional SSFT (thin solid), and the numerically obtained distribution function $P_\tau(pw\tau)$, scaled by its value at zero (thin dashed).

3.2 Large deviation method – extended SSFT

Therefore, we used next an analytical asymptotic approach based on large deviation theory[14] similar to the treatment by Lebowitz and Spohn[5]. One considers then

$$e(\lambda) \equiv \lim_{\tau \rightarrow \infty} -\frac{1}{w\tau} \ln \langle e^{-\lambda Q_\tau} \rangle. \quad (12)$$

This infinite- τ quantity is used to reconstruct the distribution function of Q_τ for large τ by setting

$$P_\tau(Q_\tau) \sim \exp [-w\tau \hat{e}(Q_\tau/w\tau)], \quad (13)$$

where $\hat{e}(p)$ is the Legendre transform of $e(\lambda)$:

$$\hat{e}(p) = \max_{\lambda} [e(\lambda) - \lambda p]. \quad (14)$$

For a class of models, Lebowitz and Spohn proved the symmetry relation

$$e(\lambda) = e(1 - \lambda). \quad (15)$$

From this, using equations (11), (13) and (14), one sees that $\lim_{\tau \rightarrow \infty} f_\tau(p) = p$, i.e., the conventional SSFT holds[5].

Our numerical results suggest, however, that for our model the conventional SSFT for the heat does not hold. We therefore expect equation (15) to be violated. Indeed, the following calculation of $e(\lambda)$ shows this to be the case.

The Fourier transform $\hat{P}_\tau(q)$ in equation (10) determines $e(\lambda)$. First, from equation (8), we have

$$\langle e^{-\lambda Q_\tau} \rangle \equiv \int_{-\infty}^{\infty} dQ e^{-\lambda Q_\tau} P_\tau(Q_\tau) = \hat{P}_\tau(i\lambda), \quad (16)$$

Thus we need the analytic continuation of $\hat{P}_\tau$ to imaginary arguments. This poses no difficulty as long as $\hat{P}_\tau$ remains analytic. One finds from equations (10) and (16)

$$\langle e^{-\lambda Q_\tau} \rangle = \frac{\exp \left[-w\lambda(1 - \lambda) \left\{ \tau + \frac{2\lambda^2(1 - e^{-\tau})^2}{1 - (1 - e^{-2\tau})\lambda^2} \right\} \right]}{[1 - (1 - e^{-2\tau})\lambda^2]^{3/2}}. \quad (17)$$

Clearly, there are divergences at the singular points $\lambda = \pm(1 - e^{-2\tau})^{-1/2}$, where the right-hand side (r.h.s.) of equation (16) is no longer analytic, so that equation (17) only holds for values of λ in between those. Using equations (12) and (17), we have

$$e(\lambda) = \lambda(1 - \lambda) \quad \text{for } |\lambda| < 1, \quad (18)$$

where, taking $\tau \rightarrow \infty$ as in equation (12) moves the singularities to ± 1 . This $e(\lambda)$ satisfies equation (15) for $0 < \lambda < 1$.

However, as λ approaches the singularities, the function in equation (17) diverges. Beyond the singularities at $\pm(1 - e^{-2\tau})^{-1/2}$, the r.h.s. of equation (17) becomes purely *imaginary*, and multivalued due to the denominator. But the *left-hand side* of equation (17) remains *real*. Clearly, we cannot use equation (17) for $|\lambda| > (1 - e^{-2\tau})^{-1/2}$. To determine $\langle e^{-\lambda Q} \rangle$ in that case, we first need to know why the integral in equation (16) diverges as $\lambda \rightarrow \pm(1 - e^{-2\tau})^{-1/2}$. As we will argue next, this happens because P_τ

has exponential tails. Since $P_\tau(Q_\tau)$ is a normalized distribution and $e^{-\lambda Q_\tau}$ a regular function, the divergence in equation (16) can only be due to the behavior of the integrand at $\pm\infty$. In fact, for $\lambda > 0$, any divergence must be due to the behavior at negative Q_τ and for $\lambda < 0$, it must be due to the behavior at positive Q_τ . Now, for $\lambda < 0$, if the distribution function $P_\tau(Q_\tau)$ fell off faster than exponential for large positive Q_τ , the factor $e^{-\lambda Q_\tau}$ could not make the integral diverge. But as it does diverge, we conclude that the distribution function falls off exponentially or slower. On the other hand, if it did fall off slower than exponential, then the exponential factor $e^{-\lambda Q_\tau}$ would always dominate the distribution function for large positive Q_τ and the integral in equation (16) would diverge for all $\lambda < 0$. Since there *are* negative values of λ for which the integral converges, the function P_τ cannot fall off slower than exponential. Hence, it must fall off exponential for large Q_τ . Considering $\lambda > 0$, one deduces along similar lines that it must fall off exponentially for large negative values of Q_τ as well.

In fact, the integral in (16) diverges for all $|\lambda| \geq (1 - e^{-2\tau})^{-1/2}$. For if the function $P_\tau(Q_\tau)$ falls off exponentially for large positive Q_τ , say as e^{-aQ_τ} , the integral in (16) diverges for all $\lambda \leq -a$. Likewise, given that the $P_\tau(Q_\tau)$ falls off like e^{aQ_τ} for large negative Q_τ , the integral diverges for all $\lambda \geq a$. Hence, for $|\lambda| \geq (1 - e^{-2\tau})^{-1/2}$, the quantity on the r.h.s of equation (12) of which the limit is taken, is minus infinity for all τ , so that $e(\lambda) = -\infty$. Thus, equation (18) becomes

$$e(\lambda) = \begin{cases} \lambda(1 - \lambda) & \text{for } |\lambda| < 1 \\ -\infty & \text{otherwise.} \end{cases} \quad (19)$$

This $e(\lambda)$ does *not* satisfy the symmetry relation in equation (15), e.g., for $\lambda = -1/2$, $e(\lambda) = -3/4$ whereas $e(1 - \lambda) = -\infty$. The fact that equation (15) is not satisfied, means that the conventional SSFT does not hold. To make this more precise, we use equations (14) and (19), to find

$$\hat{e}(p) = \begin{cases} -p & \text{for } p < -1 \\ (p - 1)^2/4 & \text{for } -1 \leq p \leq 3 \\ p - 2 & \text{for } p > 3. \end{cases} \quad (20)$$

Note that via equation (13), the large $|p|$ behavior is indeed exponential⁴.

⁴Singularities affected the distribution of a different quantity as well in J. Farago, J. Stat. Phys. **107**, 781 (2002).

Using equations (11), (13) and (20), we find

$$\lim_{\tau \rightarrow \infty} f_\tau(p) = \begin{cases} p & \text{for } 0 \leq p < 1 \\ p - (p-1)^2/4 & \text{for } 1 \leq p < 3 \\ 2 & \text{for } p \geq 3. \end{cases} \quad (21)$$

For negative p , we have

$$f_\tau(-p) = -f_\tau(p). \quad (21')$$

Equation (21) is an extension of the conventional SSFT. It coincides with it for the middle region $-1 < p < 1$,⁵ but differs from it for other p values. Most notably, for $p \geq 3$, it attains a constant value of two.

3.3 Saddle-point method – large finite times

If we compare the exact prediction of equation (21) (plotted as the bold solid line) with the numerical results (bold dashed line) in figure 1, a clear discrepancy emerges: The curve of f_τ keeps increasing with increasing p , whereas equation (21) predicts that it should level off to a value of two. This turns out to be a finite τ effect. To prove this, we need a better treatment for large but not infinite τ . This can be obtained from a saddle-point method applied to $e(\lambda)$, which we will present in a future publication[15]. The saddle-point method gives reliable results for sufficiently large τ , as can be verified by a comparison to our numerical results[15]. The asymptotic behavior for large τ is then given by

$$f_\tau(p) = \begin{cases} p + h(p)/\tau + O(\tau^{-2}) & \text{for } p < 1 \\ 2 + \sqrt{8(p-3)}/\tau + O(\tau^{-1}) & \text{for } p > 3, \end{cases} \quad (22)$$

where

$$h(p) = \frac{8p}{9-p^2} - \frac{3}{2w} \ln \left[\frac{(3-p)(1+p)}{(3+p)(1-p)} \right].$$

We left out the behavior in between $p = 1$ and $p = 3$ as it does not appear to give any additional insight. Equation (22) shows that as a function of p , the function $f_\tau(p)$ increases like $\sim \sqrt{p-3}$ (for fixed τ), while as a function of τ , it decreases (for fixed p). As expected, for $\tau \rightarrow \infty$, it approaches the large deviation result in equation (21) as $1/\tau$ for small p and as $1/\sqrt{\tau}$ for large p .

⁵A restriction to the SSFT (i.e., $|p| < 1$) was also found in a model of heat conduction in reference [9].

Whether the new features beyond $p = 1$ are observable depends on the values of $P_\tau(pw\tau)$ and $P_\tau(-pw\tau)$. If these are too small, the corresponding $f_\tau(p)$ cannot be seen in an experiment. But the value of the distribution function as plotted in figure 1 is non-negligible for values for p (and $-p$) at which f_τ bends away from the conventional SSFT. Furthermore, the values $v^* = 1.5$ and $\tau = 1.3$ used in figure 1 are realistic, as in the experiment of Wang *et al.* $v^* \approx 2.5$ and τ goes up to ≈ 6 . So this behavior should be experimentally detectable.

4 Discussion

Summarizing, we have shown that the behavior of heat fluctuations in a dissipative system with a *deterministic, mechanical* component (the potential), and a *stochastic* component (the heat bath, i.e., the water), differs from that known from both purely deterministic systems and purely stochastic systems, in two respects. a) For infinite τ , the behavior of the conventional SSFT is only seen for the scaled heat fluctuation p between -1 and 1 . For $p > 1$, after a parabolic region between $p = 1$ and 3 , the quantity f_τ no longer increases, but stays at a plateau value of two [for $p < -1$, see equation (21')]. b) The finite τ behavior of the conventional SSFT is in general unknown, but in our case we find that f_τ keeps increasing with p . However $f_\tau(p)$ stays well below the conventional SSFT, implying a larger ratio of the probabilities of the particle to absorb rather than supply heat. These features are observable.

One of the striking features of the extended SSFT, the plateau value of two for large (infinite) τ and large p , can be understood physically. For large τ and large Q_τ , i.e. p , the exponentially distributed ΔU_τ far outweighs the Gaussian distributed W_τ in equation (6). The distribution of ΔU_τ is exponential for large values ($\propto e^{-\beta|\Delta U_\tau|}$), because it is the difference of the potential energies of the particle at two times [cf. equation (7)], which are both Boltzmann-like distributed (because of the presence of the water) and independent of each other for large τ . As $\langle Q_\tau \rangle = w\tau$, this leads to $P_\tau(Q_\tau) \propto e^{-\beta|Q_\tau - w\tau|}$, which yields $P_\tau(Q_\tau)/P_\tau(-Q_\tau) \approx e^{2\beta w\tau}$ (if $Q_\tau > 0$). Then equation (11) and $\beta = 1$ give $f_\tau \approx 2$ of equation (21).

We only considered an extension of the SSFT here. An extension of the TFT can also be obtained. While for $\tau \rightarrow \infty$, one gets again equation (21), for finite times, the extended TFT and SSFT differ[15]. Furthermore, the extended TFT differs fundamentally from the conventional TFT, in that the

latter holds as an identity for all τ , whereas the former one holds only in the $\tau \rightarrow \infty$ limit[6].

Finally, one may wonder about the generality of the extended SSFT. Our Langevin-based theory is only applicable near equilibrium. The arguments given above suggest that the extended SSFT could hold for other potentials as well. Perhaps it could even hold for a larger class of systems, not near equilibrium, with mechanical and stochastic components, since these are the main physical ingredients in our theory.

Acknowledgments

We wish to thank M.J. Feigenbaum, F. Bonetto, and S. Ciliberto for useful discussions. This work was supported by the Office of Basic Engineering of the US Department of Energy, under grant No. DE-FG-02-88-ER13847.

References

- [1] D.J. Evans, E.G.D. Cohen, and G.P. Morriss, Phys. Rev. Lett. **71**, 2401 (1993).
- [2] D.J. Evans and D.J. Searles, Phys. Rev. E **50**, 1645 (1994); D.J. Searles and D.J. Evans, J. Chem. Phys. **113**, 3503 (2000).
- [3] G. Gallavotti and E.G.D. Cohen, Phys. Rev. Lett. **74**, 2694 (1995); J. Stat. Phys. **80**, 931 (1995).
- [4] J. Kurchan, J. Phys. A, Math. Gen. **31**, 3719 (1998).
- [5] J.L. Lebowitz and H. Spohn, J. Stat. Phys. **95**, 333 (1999).
- [6] E.G.D. Cohen and G. Gallavotti, J. Stat. Phys. **96**, 1343 (1999).
- [7] C. Maes, J. Stat. Phys. **95** 367 (1999); C. Maes and E. Verbitskiy, Commun. Math. Phys. **233**, 137 (2003).
- [8] G.M. Wang, E.M. Sevick, E. Mittag, D.J. Searles, and D.J. Evans, Phys. Rev. Lett. **89**, 050601 (2002).
- [9] L. Rey-Bellet and L.E. Thomas, Ann. H. Poincaré **3**, 483 (2002).

- [10] R. van Zon and E.G.D. Cohen, Phys. Rev. E **67**, 046102 (2003).
- [11] O. Mazonka and C. Jarzynski, cond-mat/9912121.
- [12] E.M. Sevick, D.R.M. Williams, D.J. Searles and D.J. Evans (private communication).
- [13] W.H. Press, S.A. Teukolsky, W.T. Vetterling, and B.P. Flannery, *Numerical Recipes in Fortran, The Art of Scientific Computing* (Cambridge University Press, Cambridge, 1992), 2nd ed.
- [14] H. Spohn, *Large Scale Dynamics of Interacting Particles*, (Springer-Verlag, Heidelberg Berlin, 1991), p. 241; *ibid.*, references ad 3.7 on p. 251.
- [15] R. van Zon and E.G.D. Cohen (2003, unpublished).